

## Evaluation of the productivity of seven varieties of wheat (*Triticum aestivum*) through integrated soil fertility management in Kaziba, South Kivu, DR Congo

Serge Shakanye Ndjadi, Bite Mubalama Mirindi, Paul Musafiri, Géant Chuma Basimine, Eloïs Cinyabuguma Lwahamire and Espoir Bisimwa Basengere

---

**Publisher**

Institut Veolia Environnement

**Electronic version**URL: <http://factsreports.revues.org/4244>

ISSN: 1867-8521

**Electronic reference**

Serge Shakanye Ndjadi, Bite Mubalama Mirindi, Paul Musafiri, Géant Chuma Basimine, Eloïs Cinyabuguma Lwahamire and Espoir Bisimwa Basengere, « Evaluation of the productivity of seven varieties of wheat (*Triticum aestivum*) through integrated soil fertility management in Kaziba, South Kivu, DR Congo », *Field Actions Science Reports* [Online], Vol. 9 | 2016, Online since 08 December 2016, connection on 09 December 2016. URL : <http://factsreports.revues.org/4244>

---

This text was automatically generated on 9 décembre 2016.

Creative Commons Attribution 3.0 License

---

# Evaluation of the productivity of seven varieties of wheat (*Triticum aestivum*) through integrated soil fertility management in Kaziba, South Kivu, DR Congo

Serge Shakanye Ndjadi, Bite Mubalama Mirindi, Paul Musafiri, Géant Chuma Basimine, Eloïs Cinyabuguma Lwahamire and Espoir Bisimwa Basengere

---

## Introduction

- 1 Every year, more than 600 million tons of wheat are harvested around the world, making wheat the second most widely cultivated cereal in the world after maize [1; 2; 3]. Wheat is the foremost cereal in terms of international trade, with 127 million tons of wheat being bought and sold in 2010 [4; 6; 7]. Wheat is a temperate climate crop, sensitive to high temperatures: for every 2°C rise in the daily temperature above 18°C, the potential yield of wheat diminishes by about 10%.
- 2 From 1996 to 1998, yields – recorded using varieties imported from France and Italy – oscillated between 200 and 400 kg/ha, depending on the agro-ecological zone (1600 to 2500 m altitude) [12; 13; 8]. In recent years the *Institut National d'Etudes et de Recherches Agronomiques* (INERA) has carried out a series of studies on wheat, conducting adaptation trials on varieties of wheat sown at their facility in order to assess the possibility of introducing varieties well adapted to the medium- and high-altitude conditions of South Kivu. Unfortunately the soils in this area are renowned for their rapid degradation and low fertility: a major constraint on agricultural production in general – and wheat-growing in particular [18; 19; 20; 21; 22].

- 3 The proposed remedy of mineral fertilization remains, in itself, fragile on soils with a high mobility of Al and Fe, which precipitate with the added nutrient elements. Also, low rural income levels in the region, coupled with the demands of applying mineral fertilizers, often represent a barrier to their acquisition. In this context, combined fertilization should offer an appropriate solution for restoring soil fertility [17; 15; 11]. Combined amendments applied to acidic tropical soils help to increase crop yields by supplying the nutrients necessary for plant growth; the improvement in the physical and biological properties of the soil enables the potential of each variety to be expressed [14; 10; 11; 16].
- 4 This paper explores the hypothesis that the seven varieties will behave differently in response to NPK, to manure and to NPK + manure. The overall aim is to evaluate the productivity of these seven varieties in response to the various treatments applied.

## Environment, materials and method

### Environment

#### Description of the environment

- 5 The experiment was conducted in a rural location at Kaziba in the territory of Walungu (1835 m, 2°48'31.6" S, 28°49'13.2" E), 55 km southwest of the city of Bukavu in South Kivu province, Democratic Republic of the Congo.
- 6 The community of Kaziba is stretched out across the high mountains of the Central African Graben to the east of the Congo River [23], at an altitude of between 1500 and 3200 m; its surface area of 192 km<sup>2</sup> is home to an average of 45,000 dwellings [24].
- 7 Its humid tropical climate, influenced by the altitude, is characterized by two seasons: the dry season, June to August, and the rainy season, September to May; mean annual temperature is about 19°C in the north and 10°C in the south, and mean annual precipitation is 1200-1700 mm. Above the 2200-m line there are mountain forests and bamboos [25].

### Materials and method

#### Description of the trial


- 8 The trial was conducted, with three replications, using the split-plot method, the main factors being the variety, with seven modalities (Farari, Kayira, Kima, Lokale, Mbega, Nyumbu and Popo) and the treatment, with four modalities (Control, NPK, Manure, and NPK + Manure).
- 9 The grains of the varieties used (obtained from INERA in Mulungu) were sown in rows 25 cm apart, with 4 grains to each pocket, every 20 cm. They were sown on the same day that the amendments were applied. Well-composted cow manure, with known characteristics, was applied to the plots, buried in the soil just before sowing at the recommended rate of 10 tons per hectare. NPK 17-17-17 was added at 150 kg per hectare. The plots marked for the NPK/manure combination received one half of each type of amendment. Upkeep consisted of regular weeding and ridging. The observations related primarily to the growth and yield parameters, centering mainly on the germination rate,

the days to flowering, days to physiological maturity, number of fertile tillers per plant, number of grains per ear, weight per 1000 grains, and yield. Variance analysis (Anova) was performed to measure differences between varieties and between treatments, as well as their interactions, and Tukey's mean separation test (at 5%) was performed using Statistix 8.0.

## Results


- 10 The germination rate was not influenced by the varieties ( $P = 0.2531$ ); nor was it affected by applications of fertilizer ( $P = 0.9980$ ) or by the interaction between variety and fertilizer ( $P = 0.3118$ ). The average germination rate was 65.04%.

Fig. 1. Effects of varieties and treatments on germination rate of wheat.




- 11 The varieties ( $P = 0.0000$ ) induced highly significant effects on the number of days to flowering. The earliest-flowering variety was Popo, after just 47 days, followed by the Farari, Kayira, Kima and Lokale varieties, which flowered after 60 days; the latest-flowering was Mbega, after Nyumbu, which flowered at 77 days and 65 days respectively.

Fig. 2. Effects of varieties and treatments on days to flowering of wheat.




- 12 The number of days to physiological maturity was correlated only to variety ( $P < 0.001$ ). The overall average for all of the varieties was 106 days; the most precocious variety was Popo (90 days) and the latest to mature was Mbega (120 days).

Fig. 3. Effects of varieties and treatments on days to physiological maturity of wheat.



- 13 The number of tillers per plant was influenced only by the variety ( $P = 0.0008$ ). Kayira had the largest average number of tillers (4.46), while Farari had relatively fewer tillers (2.15).


Fig. 4. Effects of varieties and treatments on number of fertile tillers of wheat.



- 14 Variant analysis reveals that the number of grains per ear was affected both by the variety ( $P = 0.0000$ ) and by the fertilizer applied ( $P = 0.0069$ ), but not by the crossed effects


of varieties and treatments ( $P=0.1247$ ). The variety with the most grains per ear was Lokale (43.817), while Mbega bore the least grains per ear (26.350).

Fig. 5. Effects of varieties and treatments on number of grains per ear of wheat.



- 15 The weight per 1000 grains was correlated to variety ( $P=0.0000$ ). The fertilization rate ( $P=0.3525$ ) and the interactions between these two factors ( $P=0.5629$ ) did not show an effect on the weight per 1000 grains. The overall average was 29.294 g, with Mbega and Farari at either end of the scale (at 35.56 and 22.39 grams respectively).


Fig. 6. Effects of varieties and treatments on weight per 1000 grains of wheat.



## Yield

- 16 The fertilizers show significant effects on yield ( $P=0.0194$ ); the varieties, and the interaction between the two factors, also affect this parameter ( $P=0.0013$  and  $0.0022$  respectively). The conclusion that emerges is that the plots treated with the NPK + Manure combination (1317.2 kg), produced higher yields than the unfertilized plots (605.1 B) and those treated with NPK (755.4 AB), while those treated only with manure produced values intermediate between the two (1239.6 AB). In terms of varieties, Kayira (1584.2 A) was the most productive and Mbega the least productive. As regards interactions between treatments and varieties, Manure X Kayira (2874.9 kg) and NPK + Manure X Farari (2415.0 kg) produced the best results.

Fig. 7: Effects of varieties and treatments on yield of wheat



## Discussion

- 17 The results of the variant analysis indicate that the factors analyzed, and some of their interactions, affected the parameters studied to different degrees. This demonstrates that the different varieties have different potentials, and that these potentials can be expressed more easily if the soil is amended in the appropriate way. This supports the ideas of several authors who affirm that varieties and cultivars within plant species differ in their growth and development, and that these differences are attributable to the plants' morphological, physiological and biochemical processes and their interaction with the climate, the soil, and any other practice integrated into the crop system [26; 27; 28].
- 18 According to reference [29] the average yield per hectare is 3.77 tons, too much at variance with our own results, whereas for [30], the yield is 1.92 tons per hectare, partly aligned with some of our results. The plots treated with the NPK + manure combination (1317.2 kg), produced higher yields than the unfertilized plots (605.1B), while those treated only with manure (1239.6 AB) or NPK (755.4 AB) produced yields intermediate between the two. In terms of varieties, Kayira (1584.2A) outperformed Lokale, Popo, Nyumbu and Kima, with the two remaining varieties – Farari and Mbega – ranked in an intermediate position. The crossed effects of fertilizer and variety show that the interaction Manure X Kayira (2874.9 A) produced the best results. This logical sequence in the production capacity of the varieties remains similar to that found at their center of origin [31].
- 19 These results are confirmed by [32] which states that, in general, the application of organic amendments in combination with chemical fertilizers tends to increase crop yield in comparison to plots where the same fertilizers are used in isolation; [33; 34] show that organic fertilizers have a synergetic effect with chemical fertilizers in tropical soils. The addition of mineral fertilizer – notably nitrogen and phosphorus – could increase the biological activity of the soil and, in so doing, facilitate the effective release of nutrients for the crop.
- 20 The recorded increase in yield is attributed to the improvement in the soil properties [35] and the release of nutrient elements [36]. Likewise, the increase in yield due to the input of organic amendments is attributable to the favorable change in soil conditions, leading to good root development and good assimilation of the nutrient elements released by the organic matter itself, or through retention of the nutrients released by the fertilizers [37; 38; 39].

- 21 According to [40], wheat – despite being an under-utilized crop in most tropical ecosystems, unlike other cereals such as maize and sorghum, due to its sensitivity to a range of fungal diseases, its incompatibility with the environmental conditions, and its low yield in these ecosystems compared to the latter crops – could nonetheless have a future thanks to high-performance varieties, appropriate crop systems, and well-managed soil fertility. The results obtained provide sufficient evidence to support that author's position: through use of the right variety, suitable agricultural techniques, and effective management of soil fertility, the yield of wheat was improved to a significant degree.

## Conclusion

- 22 To help promote wheat-growing in rural areas of South Kivu, DR Congo, seven varieties of wheat were evaluated using different fertilizer options: organic, based on cow manure; mineral, with NPK 17-17-17; and organo-mineral, by combining both types. The trials were conducted at Kaziba, a mountainous but densely populated region where the soil is in an advanced state of degradation. The results obtained indicate that wheat-growing is possible in this region, thanks to climate conditions that match the crop's requirements in terms of temperature. Although the soil is severely degraded, this can be remedied using organic amendments (with cow manure) and organo-mineral amendments, making the necessary nutrients available to the plant for its growth and development. Mineral fertilizers alone did not show significant effects on the crop; however, with the use of manure, and its combination with mineral fertilizers, favorable effects were observed. Of the seven varieties proposed by INERA, Kayira and Farari are the preferred options due to their high level of performance. The main recommendation is to use the variety Kayira under cow manure, providing the latter is very well composted and comes from well-fed cattle, and also to facilitate the task of improving soil fertility; combining the manure with NPK offers most additional benefits.

---

## BIBLIOGRAPHY

- [1] Laura, P. La volatilité du prix du blé sur les marchés agricoles mondiaux : une approche par les facteurs. Masters dissertation, University of Lyon; 2012.
- [2] Carême, C, Saghier, T. Conséquences de la nuisibilité des mauvaises herbes sur la production du blé d'hiver en Tunisie : les seuils d'intervention et la rentabilité du désherbage. *Tropicultura* 1991;9(2): 53-57.
- [3] Tanner, D. The maize and wheat network: catalysing for better research and impact. *Agriforum* 2000;11: 12-13.
- [4] Mushambanyi, T. Contribution à la promotion de la culture du blé (*Triticum aestivum* L.) au Sud-Kivu, RD Congo : Evaluation du potentiel de rendement de deux génotypes d'origine burundaise, dans différentes zones agro-écologiques locales. *Tropicultura* 2002;20(4): 210-216.

- [5] Sharma, RC. Analysis of phytomass yield in wheat. *Agronomy Journal* 1992;84(6): 926-929.
- [6] Bennaceur, M, Rahmoune, C, El-Jaafri, S, Opaul, R. Potentialités de production de quelques variétés de blé dur (*Triticum drums Desf.*) au Maghreb. *Rev. Sci. Technol., Univ. Constantine* 1997;8: 69-74.
- [7] Hakimi, E. Sélection sur la base physiologique et utilisation des espèces tétraploïdes du genre *Triticum* pour l'amélioration génétique de la tolérance a la sécheresse du blé. Doctoral thesis, Montpellier; 1995.
- [8] Bishweka, A. La culture du froment en Afrique Centrale (Sud-Kivu). *Recherches africaines* 1999;4: 99-112.
- [9] Gate, P. *Ecophysiologie du blé*. Ed. ITCF, Technique et Documentation, Lavoisier, Paris; 1995.
- [10] Zapata, F, Roy, RN (eds.) *Use of Phosphate Rocks for Sustainable Agriculture*. Rome, FAO; 2004.
- [11] Nyembo, KL, Kisimba, MM, Theodore, MM, Jonas, L, Kanyenga, LA, Becker, NK, Mpundu MM, Baboy LL. Effets de doses croissants des composts de fumiers de poules sur le rendement de chou de chine (*Brassica chinensis L.*). *J. Appl. Biosc.* 2014;77: 6505-6522.
- [12] Inspection provinciale d'agriculture. Rapport de l'inspection agricole de la province du Sud-Kivu, R.D. Congo. Rapport pour l'exercice de l'année 1996 à l'année 2000. Bukavu; 2000.
- [13] Payne, T. Impact of maize and wheat research in Eastern and Central Africa: Results of recent studies. *ASARECA/Agriforum* 2000;10: 7-9.
- [14] Kotchi, V, Yao, KA, Sitapha, D. Réponse de cinq variétés de riz à l'apport de phosphate naturel de Tilmesi (Mali) sur les sols acides de la région forestière de Man (Côte d'Ivoire). *J. Appl. Biosc.* 2010;31: 1895-1905.
- [15] Mulaji, KC. Utilisation des composts de biodéchets ménagers pour l'amélioration de la fertilité des sols acides de la province de Kinshasa (République Démocratique du Congo). Doctoral thesis, Gembloux Agro bio tech; 2011.
- [16] Useni, SY, Glady, MI, Theodore, MM, Becker, NK, Jonas, L, Mick, ABL, Antoine, KL and Louis, BL. Amélioration de la qualité des sols acides de Lubumbashi (Katanga, RD Congo) par l'application de différents niveaux de compost de fumiers de poules. *J. Appl. Biosc.* 2014;77: 6523-6533.
- [17] Bulson, J, Snaydon, RW, Stopes, C. Effects of plant density on intercropped wheat and field beans in an organic farming system. *Journal of Agricultural Science* 1997;128: 59-71.
- [18] Pypers, P, Sanginga, JM, Kasereka, B, Walangululu, M, Vanlauwe, B. Increased productivity through integrated soil fertility management in cassava-legume intercropping systems in the highlands of Sud-Kivu, DR Congo. *Field Crops Research* 2011;120: 76-85.
- [19] Vanlauwe, B, Mutuo, P, Mahungu, N, Pypers, P. Boosting the productivity of cassava-based systems in DR Congo. IITA. R4D Review 9, by B. Vanlauwe & K. Lopez, eds. Ibadan, Nigeria; 2012: 30-34.
- [20] Bagula, M. Evaluation de l'efficacité d'usage des engrais dans les sols dégradés du Sud-Kivu sur la culture du maïs et du haricot. Cas du groupement de Burhale. Engineering degree dissertation, UEA, unpublished; 2010.
- [21] World Bank, Africa Region. *Promoting Increased Fertilizer Use in Africa: Lessons Learned and Good Practice Guidelines*. Africa Fertilizer Strategy Assessment, ESW Technical Report, Washington, USA; 2006.

- [22] IFDC. Rapport des activités effectuées à l'est de la République Démocratique du Congo, Kigali; 2009.
- [23] Mulumeoderhwa, L. Elevage de bovins dans la collectivité de Kaziba, ISP / Bukavu. TFC, unpublished; 1995.
- [24] Bashagaluke, J. Analyse diagnostique des systèmes de culture à base de manioc et la gestion de l'érosion au Bushi. Engineering degree dissertation, U.C.B., unpublished; 2008.
- [25] Ellen, V. Nutrient deficiencies in soils of Walungu, South-Kivu, Democratic Republic of Congo. Université Catholique de Louvain; 2008.
- [26] Baligar, VC, Fageria, NK, He, ZL. Nutrient use efficiency in plants. *Communications in Soil Science and Plant Analysis* 2001;32(7-8): 921-950.
- [27] Stephenson, K, Amthor, R, Mallowa, S, Nungo R, Maziya, B, Gichuki, S, Mbanaso, A, Manary, M. Consuming cassava as a staple food places children 2-5-year old at risk for inadequate protein intake: an observational study in Kenya and Nigeria. *Nutrition Journal* 2010;9(9).
- [28] Magrini, M, Triboulet, P, Bedoussac, L. Pratiques agricoles innovantes et logistique des coopératives agricoles : Une étude ex ante sur l'acceptabilité de cultures associées blé dur-légumineuses. *Economie rurale* 2013;6(338): 25-45.
- [29] Benbelkacem, A, Kellou, K. Evaluation du progrès génétique chez quelques variétés de blé dur (*Triticum L. var durum*) cultivées en Algérie. *CIHEAM, série A*, 2000: 105-110.
- [30] Lounes, Y, Guerfi, A. Contribution à l'étude du comportement agronomique de 27 nouvelles variétés de blé dur en vue de leur inscription au catalogue officiel national. State agronomy engineering diploma, Université Mouloud Mammeri de Tizi Ouzou, Algeria; 2010.
- [31] INERA Mulungu, Annual Report 2013, Bukavu, DR Congo.
- [32] Mahadeen, A. Influence of Organic and Chemical Fertilization on Fruit Yield and Quality of Plastic-House Grown Strawberry. *Jordan Journal of Agricultural Sciences*, 2009;5(2): 167-177.
- [33] Sanginga, N, Woome, P. Integrated soil fertility management in Africa: principles, practices and process development, TSBF-CIAT and FORMAT, Nairobi, Kenya; 2009.
- [34] Bationo, A. Integrated soil fertility management option for agriculture intensification the Sudano-Sahelian zone in West Africa. Academy of Science Publishers, Nairobi, Kenya; 2008.
- [35] Togun, A, Akanbi, W. Comparative effectiveness of organic-based fertilizers to mineral fertilizers on tomato growth and fruit yield. *Compost Science and Utilization* 2003;11(4): 337-342.
- [36] Turemis, N. The effects of different organic deposits on yield and quality of strawberry cultivar Dorit. *Acta Hort.* 2002;567: 507-510.
- [37] Amanullah, M, Alagesan, A, Vaiyapuri, K, Sathyamoorthi, K, Pazhanivelan, S. Effect of intercropping and organic manures on weed control and performance of cassava (*Manihot esculenta* Crantz). *Journal of Agronomy* 2006;5(4): 589-594.
- [38] Alley, M, Vanlauwe, B. The role of fertilizer in integrated plant nutrient management. IFA and TSBF-CIAT, Paris; 2009.
- [39] Compère, R. Maintien et restauration de la fertilité des sols en région sahélo soudanienne sénégalaise par une association rationnelle des activités d'élevage et d'agriculture. *Bulletin Recherche Agronomique* 1991;26: 153-167.
- [40] Gleizes, J. Des chiffres et des céréales : l'essentiel de la filière. *Passion céréales*, Paris; 2011.

## ABSTRACTS

Due to high demand for food resources as a result of increasing population, the promotion of large-scale crops such as wheat has become essential. Unfortunately, soil infertility and a lack of improved seed are major constraints on the expansion of this crop in Kaziba, a mountainous rural area in South Kivu. The productivity of seven wheat varieties (Farari, Kayira, Kima, Lokale, Mbega, Nyumbu and Popo) was evaluated under organic and mineral fertilizer during the 2013-2014 crop season on poor soil in South Kivu (DR Congo). NPK 17-17-17, farmyard manure and their combination were applied as fertilizers in a split-plot trial design with three replications. The observations focused on the growth and yield parameters, and the results revealed differences between varieties, treatments and interactions. The NPK + farmyard manure treatment gave the highest mean yield (1317.2 kg), Kayira was the most productive variety (1584.2 kg), and the interaction Kayira X farmyard manure was the most effective (2874.9 kg). The variety Kayira would seem to be indicated, with farmyard manure as the recommended fertilizer, being locally accessible and easily usable for promoting wheat in the region.

## INDEX

**Keywords:** wheat, productivity, varieties, poor soil, fertilizers

## AUTHORS

### SERGE SHAKANYE NDJADI

Masters in sustainable soil management. *Université Evangélique en Afrique*, Faculty of Agricultural and Environmental Sciences, Crop production Department

### BITE MUBALAMA MIRINDI

Engineer in plant production, *Université Evangélique en Afrique*, Faculty of Agricultural and Environmental Sciences, Crop production Department

### PAUL MUSAFIRI

Masters in soil science, Jomo Kenyatta University, Soil Science Department

### GÉANT CHUMA BASIMINE

Engineer in plant production, *Université Evangélique en Afrique*, Faculty of Agricultural and Environmental Sciences, Crop production Department

### ELOÏS CINYABUGUMA LWAHAMIRE

Masters in tropical and subtropical plants protection), *Institut National d'Etudes et de Recherches Agronomiques*, National Cereals Program

### ESPOIR BISIMWA BASENGERE

PhD in agricultural science, Catholic University of Bukavu, Faculty of Agricultural and Environmental Sciences, Crop production Department